

Reliability of sleep deprivation-associated spontaneous brain activity and behavior

Lei Gao^{1,2†}, Lijun Bai^{3†}, Yuchen Zhang⁴, Xi-jian Dai², Rana Netra¹, Youjiang Min⁵, Fuqing Zhou², Honghan Gong^{2*}, Ming Zhang^{1*}, and Yijun Liu^{6,7}

1. Department of Medical Imaging, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, China

2. Department of Radiology, the First Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

3. The Key Laboratory of Biomedical Information Engineering, Ministry of Education, Department of Biomedical Engineering, School of Life Science and Technology, Xi'an Jiaotong University, Xi'an 710049, China

4. Zonglian Experimental Class, Xi'an Jiaotong University, Xi'an 710049, China

5. Acupuncture & Rehabilitation Department, Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine, Nanchang 330006, Jiangxi Province, China

6. Southwest University, School of Psychology, Key Laboratory of Cognition and Personality, Chongqing, Chongqing 400715, China

7. Departments of Psychiatry and Neuroscience, McKnight Brain Institute, University of Florida, Gainesville, FL 32610, USA.

* Correspondence to Honghan Gong, Department of Radiology at the First Affiliated Hospital of Nanchang University, 17 Yongwaizheng Ave., Nanchang 330006, Jiangxi Province, China
Email: honghan_gong@sina.com (Hong-Han Gong);

and Ming Zhang, Department of Medical Imaging at the First Affiliated Hospital of Xi'an Jiaotong University, 277 West Yanta Road, Xi'an 710061, Shaanxi Province, China. Email: zmmri@163.com (Ming Zhang)

† Contribute equally to this paper.

Abstract

Recent studies have indicated that sleep deprivation (SD) alters intrinsic low-frequency connectivity in the resting brain, mainly focusing on the default mode network (DMN) and its anticorrelated network (ACN). These networks hold key functions in segregating internally and externally directed awareness. However, far less attention has been paid to investigation of the altered amplitude of these low-frequency fluctuations (ALFF) at the whole-brain level and more importantly by what extent the sleep-deprived resting brain pattern can be reproducible and predict individual behavioral performance. The aim of this study was to characterize more clearly the influence of sleep on the whole brain level of ALFF changes and its relation with the performance of a lexical decision task in the sleep deprivation. Sixteen healthy participants underwent fMRI three times: once after a normal night of sleep in the rested wakefulness (RW) state and two following approximately 24 h of total SD separated by an interval of two weeks (SD1 and SD2). Our behavioral results showed that sleep stabilizes performance whereas two sleep deprivation even at an interval of two weeks consistently deteriorates it. Sleep deprivation attenuated the ALFF mainly in the bilateral orbitofrontal cortex (OFC), bilateral dorsolateral prefrontal cortex (DLPFC) and right inferior parietal lobule (IPL). By contrast, the enhanced ALFF emerged in the left sensorimotor cortex (SMA), visual cortex and left fusiform gyrus. Conjunction analysis of SD1 and SD2 versus the control maps and voxel-wise ICC analysis revealed that these SD induced ALFF changes showed a significantly high reliability ($ICC > 0.5$). Particularly, the attenuation of the right IPL presents a significant negative relation with the behavior performance and can be reproducible for two SD at an interval of two weeks. Our results suggest that ALFF is a stable measure in study of SD, and the right IPL may represent a stable biomarker that responds to sleep loss.

Keywords: sleep deprivation, resting-state fMRI, ALFF, test-retest, reliability

Introduction

Sleep deprivation (SD) repeatedly show a variable (negative) impact on mood, cognitive performance, and motor function due to increasing sleep propensity and destabilization of the wake state (Goel et al., 2009). Previous neuroimaging studies have shown that insufficient sleep can have a number of adverse effects on brain functioning. The investigation of the neural mechanism underlying SD by using fMRI is still in the initial stage, but has already provided a wealth of information about sleep-deprived brain networks and various functions that support normal waking behavior. Various neuroimaging studies have found altered activation patterns of sleep-deprived brain during a number of tasks including memory (Chee and Choo, 2004; Chee et al., 2006; Mu et al., 2005; Mu et al., 2005; Sterpenich et al., 2009; Van Dongen, 2005), attention (Chee and Tan, 2010; Chee et al., 2008; Kong et al., 2012), executive functioning (Drummond and Brown, 2001; Muto et al., 2012), and decision making (Chee et al., 2010; Kong et al., 2011; Libedinsky et al., 2011). Several brain regions, such as the prefrontal cortex, parietal cortex, sensorimotor area, visual cortex, thalamus and cingulate cortex have been frequently reported to be associated with SD.

Prolonged wakefulness has been associated with altered functional integrations in the resting brain. The default mode network (DMN), with a relatively high rCBF (Gusnard et al., 2001; Raichle et al., 2001) and a high level of correlated BOLD signal fluctuations (Greicius et al., 2003; Raichle, 2011) at rest, is considered to support self-awareness (Gusnard et al., 2001) and conscious self-representation (Lou et al., 2004). Investigations of the influence of SD on the DMN had shown significantly disrupted deactivations and led to double dissociations within anterior as well as posterior midline regions of the DMN (Gujar et al., 2010). This finding invites the speculation that the decreased DMN connectivity could be intrinsic to the sleep deprivation or a reflection of changes in vigilance (De Havas et al., 2012; Samann et al., 2010). In addition, the DMN is anticorrelated with the cognitive control network (CCN), a corresponding task-positive network, which encompasses bilateral fronto-cingulo-parietal structures including lateral prefrontal and superior parietal areas (Niendam et al., 2012). Abnormal connectivity within DMN and its anticorrelated networks (CCN) have also been reduced after SD (Bosch et al., 2013; De Havas et al., 2012; Samann et al., 2010; Shao et al., 2013). DMN-CCN interactions

may be considered to reflect the level of consciousness that is required for information integration (Heine et al., 2012; Larson-Prior et al., 2011). More recently, two studies pointed out that enhanced functional connectivity between the dorsal nexus and dorsolateral prefrontal cortex, as well as attenuated functional connectivity within thalamocortical, also occurs after a SD (Bosch et al., 2013; Shao et al., 2013). These findings indicate disrupted temporal synchronization of the global resting-state network in the sleep-deprived brain. However, these investigations generally adopted the seed-based functional connectivity method and did not allow for a direct assessment of regional activity during the resting brain. In other words, abnormal functional interactions between two remote areas cannot address the question on which area is responsible for such observable connectivity alternations.

In light of this view, we aimed to measure the amplitude of low-frequency fluctuation (ALFF) in sleep-deprived resting brain. ALFF, without a priori selection of regions of interest, can be used to study the dynamics of the BOLD signal at the local, voxel-wise level, without assessing the relationship between regions (Zang et al., 2007). In addition, such method has proven to be test-retest reliability across time (Li et al., 2012; Turner et al., 2012; Yan et al., 2013; Zuo et al., 2010), and can successfully predict magnitudes of the task-evoked activity (Mennes et al., 2011; Zou et al., 2013). The present study attempted to address the following issues: *i*) by which way sleep-deprived brain reflects abnormal patterns of ALFF; *ii*) whether such patterns are reproducible over a two-week interval SD; *iii*) to what extent the inter-subject differences in the ALFF resting activity may predict individual behavioral performance. For this purpose, a lexical decision task was also used as a paradigm to test the effects of sleep loss and fatigue on the dynamic time-course of responses to cognitive load (Babkoff et al., 1985; Forster and Forster, 2003; L O Pez Zunini et al., 2014).

Materials and Methods

Subjects

Sixteen healthy volunteers (8 females, mean age of 22.1 ± 0.8 years) were recruited in this study after giving the informed consent. Participants were selected from respondents to a web-based questionnaire. They should meet the following criteria: (1) right-handed according to

the modified Edinburgh Handedness Questionnaire (Oldfield, 1971); (2) between 20 and 24 years of age; (3) good sleeping habits (sleeping no less than 6.5 h each night for the past one month); (4) not be of extreme morning or evening chronotype (score no greater than 22 on a modified Morningness–Eveningness scale; Horne and Ostberg, 1976); (5) no long-term medications; (6) no symptoms associated with sleep disorders; (7) no history of any psychiatric or neurologic disorders; (8) no history of drug abuse and current use of anti-depressant or hypnotic medications. Participants had an average of 15.7 ± 1.2 years of education. This study was approved by the medical research ethics committee and institutional review board of The First Affiliated Hospital of Nanchang University.

Participants showed normal sleep quality as assessed using the Pittsburgh Sleep Quality Index (PSQI) [37] (mean \pm SD, 1.5 ± 0.97) and normal daytime sleepiness as assessed using the Epworth Sleepiness Scale (ESS)[38] (mean \pm SD, 6.44 ± 2.07). They had a BMI (in kg/m²) of 17.5–22, and were free of nightshift work. Before experiment (approximately 4 weeks before), subjects were required to sleep 7–9 h/night, preceding 00:10 a.m on average and keep sleep logs.

Sleep deprivation and experimental protocol

All subjects were scheduled for three fMRI scans starting at 7:00 PM. One scan follows the individual's normal sleep (RW group), while the other two after a night of total SD (i.e., SD1 and SD2 groups). Two SD sessions were approximately two weeks apart. The sequence of experiments was counterbalanced across sessions and approximately two weeks apart between two experiments. This was to ensure minimizing the possibility of residual effects of SD on cognition in participants whose SD session preceded their rested wakefulness session. Subjects were forbidden to tea, coffee or caffeine content drinks and alcohol intake for 72 h before fMRI examinations. Sleep logs were kept for a week prior to the study night. During SD, subjects were monitored in the lab onward and only allowed to engage in non-strenuous activities such as reading and watching videos. Vigorous physical activity prior to the scans was forbidden.

A behavior test was performed for each subject prior to fMRI scanning. Word stimuli were displayed using the experimental software DMDX v.3.0.4 (Forster and Forster, 2003). White-color words of 10 mm in size were presented on monitor with a black background in a

dimly lit room. During the experiment, each pair of stimuli was presented for 900 ms separated by a blank screen for 500 ms. Subjects had 2,500 ms between trials to judge whether the two words were semantically related or not. A positive response was indicated by pressing the right button using the middle finger of the right hand, while a negative response was represented corresponding left button to the index finger of the right hand. Participants were encouraged to proceed as quickly and accurately as possible. Accuracy and reaction times (RTs) (to the nearest millisecond) were recorded. Before the formal test, a short period of practice with a different set of sentences was provided.

Data acquisition

fMRI data were collected on a SIEMENS Trio 3.0 T scanner. Each subject lied on supine with the head in neutral position fixed comfortably by a belt and foam pads during the test. The scanning sessions included: (1) localizer, (2) T1 MPRAGE anatomy (176 sagittal slices, thickness/gap = 1.0/0 mm, in-plane resolution = 256×256 , FOV (field of view) = $240 \text{ mm} \times 240 \text{ mm}$, TR (repetition time) = 1,900 ms, TE (echo time) = 2.26 ms, flip angle = 15°), (3) EPI-BOLD (36 axial slices, echo-planar imaging pulse sequence, thickness/gap = 5.0/1 mm, in-plane resolution = 64×64 , TR = 3,000 ms, TE = 30 ms, flip angle = 90° , FOV = $240 \text{ mm} \times 240 \text{ mm}$). During the resting-state fMRI session, subjects were required to be as calm as possible, to keep their eyes closed but not to fall asleep to ensure a successful image acquisition.

Behavioral analysis

To examine changes in behavior performance over the course of the SD sessions, the mean RT and false rates (button pressed following a cue) to measure the accuracy of performance were computed respectively.

Data preprocessing

All preprocessing was performed using the Data Processing Assistant for Resting-State fMRI (DPARSF, Yan and Zang, 2010, <http://www.restfmri.net>), which is based on Statistical Parametric Mapping (SPM8) (<http://www.fil.ion.ucl.ac.uk/spm>) and Resting-State fMRI Data Analysis Toolkit (REST, Song et al., 2011, <http://www.restfmri.net>). For the resting-state fMRI

data on each subject, the first two volumes were discarded to avoid the possible effects of scanner instability and adaptation of subjects to the circumstances.

The following sequence of preprocessing steps was performed: *i*) slice timing (correction for the within-scan acquisition time differences between slices); *ii*) head motion correction (realignment and a six-parameter spatial transformation). Recent studies indicate that head motion can significantly influence measures and results derived from the resting-state fMRI (Power et al., 2012; Van Dijk et al., 2012; Yan et al., 2013), we computed the voxel-specific head motion, including voxel-specific framewise displacement (FD_{vox}) and voxel-specific total displacement (TD_{vox}) values for each subjects by using the DPARSF toolbox. Group differences of mean FD_{vox} were calculated by using two-sample t-test while with no significant group differences. Then the mean FD was used as a covariate in the group comparisons of ALFF. In our study, absolute head movement was below 0.5 mm and 0.5° for all subjects; *iii*) spatial normalization to the Montreal Neurological Institute (MNI) template (resampling voxel size = $3 \times 3 \times 3 \text{ mm}^3$); *iv*) spatial smoothing (full width at half maximum (FWHM) = 6 mm Gaussian kernel); *v*) linear detrend and voxel-wise bandpass filtering (0.01 - 0.08 Hz).

For ALFF analysis, for a given voxel, the time series was first converted to the frequency domain (0.01-0.1Hz) using a Fast Fourier Transform (FFT). The square root of the power spectrum was computed and then averaged across a predefined frequency interval. This averaged square root was termed ALFF at the given voxel. ALFF measures the absolute strength or intensity of LFF. ALFF of each voxel was computed for each participant and was further divided by the global mean value to reduce the global effect of variability across participants.

Test-retest reliability analyses

Test-retest reliable measurements are important for the inference of convincing conclusions. To investigate the test-retest reliability of SD, we further calculated voxel-wise intraclass correlation coefficients (ICC) between two SD sessions (Zuo et al., 2010).

Statistical analyses

We used paired *t*-test for SD vs. RW groups to determine the effects of SD on performance measures of RT. For ALFF, a one-sample one-sided *t*-test was performed inter-group to determine whether the ALFF differed from 1 (Raichle et al., 2001; Zang et al., 2007), and a paired *t*-test to see the differences between groups. Voxels with a *p* value < 0.01, cluster size > 1,053 mm³ (39 voxels) were considered to be a significant statistical difference between two groups, which also corresponds to a multiple corrected *p*<0.05 in the AlphaSim program (<http://afni.nih.gov/afni/docpdf/AlphaSim.pdf>).

Conjunction analysis

A conjunction of SD1 and SD2 versus control maps was calculated to determine brain areas that commonly and similarly differently in both SD1 and SD2.

Brain-behavior relationships

To further evaluate the relationship between ALFF changes and behavioral performance after SD, we examined the Pearson's correlation between the mean ALFF values of peak voxels of areas derived from the inter-group comparison and behavioral performance of RT.

Results

Performance findings

There were no significant differences in the accuracy rate between the SD and RW groups. The two SD groups had significantly longer reaction time than that of the RW group (RW = 2,010.4 ± 227.17 (ms), SD1 = 2,275.1 ± 176.66 (ms), *T* = -3.858, d.f. = 15, *P* < 0.002; SD2 = 2,172.2 ± 166.51 (ms), *T* = -2.584, *p* < 0.021) (Fig. 1).

ALFF differences in two SD sessions

Comparing with the RW group, both SD groups showed the decreases of ALFF mainly in the bilateral orbitofrontal cortex (OFC) (BA 11/47), bilateral dorsolateral prefrontal cortex (DLPFC) (BA 46) and right inferior parietal lobule (IPL, BA39/40), while the increases of ALFF

emerged primarily in the somatosensory cortex (SMC), fusiform gyrus (BA 37) and middle occipital gyrus (MOG) (BA 18/19) (Fig. 2 and Table.1).

Conjunction analysis and voxel-wise ICC analysis

The overlap of SD1 and SD2, compared to control group, shared the same areas primarily in the Pcu/PCC, OFC, SMC, bilateral occipital cortex, and right IPL (shown in Fig. 3.). The high reliability derived from the intra-subject test-retest analysis existed in most of brain areas (ICC ≥ 0.5 , Fig.3). These areas mainly located at the precuneus /PCC, cingulate cortex, bilateral SMC, bilateral parietal cortex, bilateral MPFC, and bilateral occipital cortex.

Correlations between ALFF and behavior performance in the SD group

In this study, the greater the prolongations in the RT from rest to SD, the greater the decrease in ALFF in the right IPL reproducible by two SD at an interval of two weeks (for SD1, $r = -0.561$, $p < 0.024$; while for SD2, $r = -0.499$, $p < 0.05$) (Fig.4).

Discussion

Several observations emerged from the examination of spontaneous “low frequency” fluctuation from temporal synchronization (i.e. functional connectivity) in sleep-deprived resting brain. Beyond verifying the altered intrinsic patterns in “off-line” brain, we noted that even one night of sleep loss can alter the ALFF, especially attenuations in the bilateral OFC, bilateral DLPFC, and right IPL, while enhancements in the SMC, thalamus, fusiform gyrus and visual cortices. Moreover, the measurements of the ALFF patterns in the sleep-deprived brain are quite reliable over time with an interval of two weeks. These relatively high stability coefficients provided conservative estimates of the test-retest reliability of ALFF in SD. In addition, the inter-subject differences in ALFF measures provide a clue to predict the individual behavioral performance, which showed a significantly negative relation of RT with the ALFF of the right IPL.

Significantly different patterns of ALFF were found between SD and RW. Consistent with previous task-evoked activations in varied cognitive stimuli, our findings showed that altered

ALFF changes were mainly located in the DLPFC, OFC and IPL. It is believed that the frontal-parietal areas are particularly vulnerable to SD (Bosch et al., 2013; Chee and Tan, 2010; Chuah et al., 2010; Lythe et al., 2012). Decreased activities in the prefrontal and parietal were especially found in several working memory studies, therefore parietal activity represents a biomarker of individual response to sleep debt (De Havas et al., 2012). Inadequate sleep is also associated with exaggerated emotional responses (Chuah et al., 2010; Goldstein et al., 2013; Killgore, 2013; Menz et al., 2012; Minkel et al., 2012; Mullin et al., 2013). The OFC, known as the limbic system, is thought to be critical for mediating the interactions between emotional processes and cognitive functions, such as decision making (Azzi et al., 2012; Kahnt et al., 2012; Parsons et al., 2013; Zald et al., 2012). We speculate that SD may influence emotion and cognition regulation circuitry, and lead to deficient capacity to regulate emotional arousal and cognitive loads.

Besides impairments in sleep-deprived resting brain, it was also worthy to note that enhanced ALFF mainly emerged in the SMC, fusiform gyrus and visual cortex, which may reflect compensation mechanism after sleep loss, and an attempt to maintain hyperarousal. Early PET and fMRI studies investigate brain responses to attention tasks following SD, and also find such hypermetabolism in the visual cortex and sensorimotor areas (Thomas et al., 2000). They inferred that such an enhancement of brain activity might indicate a homeostatic drive for recovery of brain areas involved in attention and higher-order cognitive processes, interpreted as compensatory mechanisms to maintain alertness and cognitive performance in the face of extended wakefulness. A recent study of enhanced functional connectivity between primary sensory processing and motor planning regions in insomnia provides further evidence, consistent with our results (Killgore et al., 2013). This may imply ALFF encodes tendency for task response even during the resting state.

As expected, we found highly reproducible ALFF patterns at a two-week interval in the sleep-deprived brain. The overlapped areas presented in our study included the PCC, OFC, IPL, SMC, visual cortex and right insula via the conjunction analysis. Voxel-wise ICC map showed the spatial distribution of the reliability of the observed ALFF, mainly along the midline

structure of the brain, lateral prefrontal and parietal cortex ($p < 0.001$). This finding was in line with previous task-fMRI research (Lim et al., 2007), which has shown reproducible brain activations highly correlated across sessions in a frontoparietal network during a working memory task. As in previous studies, stable drop in the left parietal activation over time, we denoted that these tasks-evoked brain activity and behavior changes may underlie intrinsic brain activity. The reliability of the resting spontaneous activity pattern, acting as a functional framework for a moment-to-moment responses, remains uncovered. In the present study, we confirmed its stability over a two-week interval. Along the same lines, some studies also found reliability of the ALFF in chronic schizophrenia (Turner et al., 2012) and healthy state (Li et al., 2012) during a short period of time. This may support the notion that, in a relatively short duration, intra-individual resting dynamics to SD load stably exists.

At resting state, the brain featured by exhibiting an intrinsic organization that includes both “task-negative” (DMN) and its anticorrelated “task-positive” networks (ACN) (Fox et al., 2005). DMN characterized by more energetic metabolic and neural activity at rest engaged in internally focused tasks, such as autobiographical memory retrieval, implicit learning, prospection, monitoring and other internally focused thought processes (Braga et al., 2013; Raichle et al., 2001; Uddin et al., 2009). Decreased ALFF signals in the PCC, bilateral OFC (BA 11/47) and right inferior parietal lobule (IPL, BA39/40) following 24h SD may be associated with maladaptive ability to switch the default mode activity when task conditions required attentions. The DLPFC, one of the “task-positive” regions (ACN), has been implicated to be responsible for the failure in working memory in SD (Goel et al., 2009). In our study, reduced DLPFC activity may suggest dysfunctional integrations between the DMN and ACN, and this is consistent with reduced anti-correlation between DMN and ACN nodes during both task and resting states (De Havas et al., 2012). Moreover, the interesting finding is the increased ALFF in the SMC, visual cortex and left FG. According to the sleep homeostatic hypothesis (Born and Feld, 2012; Tononi and Cirelli, 2003, 2006), one possible explanation may be the compensatory reallocates—the resting brain reallocate its oscillatory dynamics resources. This reconfiguration reduces the adaptation of neural resources in response to increased cognitive challenges and emotion regulations.

chinaXiv:201711.00275v1

The results obtain reproducible behavior performance following the SD at an interval of two weeks. Besides the reliability, there is a significant negative linear correlation between the ALFF decrease in the right IPL and RT when performing lexical decision task. Plentiful studies have found that sleep loss can significantly degrade performance on tasks that incorporate a working-memory component or that otherwise are thought to require contributions from frontoparietal cortical regions (Goel et al., 2009). The IPL, as an important association area of integrating sensory information, plays a prominent role in visuospatial attention. Successful performance of the semantic priming tasks demands participants to sustain mental effort, strategically control and quickly focus their attention in order to accurately register transient stimulus events, as well as rapidly compare incoming information to that maintained in working memory. In this study, the significantly negative linear correlation may suggest that spontaneous activity of the IPL was impaired with sleep loss load. Many reports, though some divergent, are associated reduced task-evoked parietal activation with a decline in behavioral performances. One recent study from resting functional connectivity also support the notion that activity changes in the IPL may reflect the early effects of SD (De Havas et al., 2012). Therefore, it is reasonable to suggest that reduced resting state intrinsic activity in the IPL may relate to declining cognitive capacity in SD.

Limitations

There are several limitations in this study. Firstly, a relatively small sample, the statistical power lowers and limited, so the results can hardly survive a strict multiple comparison correction (e.g. FDR or FWE correction). Future studies could use a larger sample size to increase the statistical power of the study. Secondly, a lack of objective assessment of both self-reported sleepiness and mood items. Further investigations should be focused on a more detailed frequency-dependent analysis.

Conclusion

Our high test-retest reliability of ALFF patterns highlighted the notion that sleep burden reshapes the low frequency dynamics in the resting brain, and this sleep-deprived brain pattern

can predict the individual behavior performances. This finding may provide much deeper insights into the understanding of SD related underlying neurological underpinnings of cognitive declines.

Conflict of Interests

There is no competing interest.

Acknowledgements

We thank Hengyi Rao (Department of Neurology, University of Pennsylvania) for the valuable discussions about the research findings. This study was supported by the National Natural Science Foundation of China (Grant No. 81371630, 81260217, and 81371530.), the Fundamental Research Funds for the Central Universities, the Beijing Nova program (grant number Z111101054511116), and the Beijing Natural Science Foundation (grant number 4122082).

References:

- Azzi, J.C., Sirigu, A., Duhamel, J.R., 2012. Modulation of value representation by social context in the primate orbitofrontal cortex. *Proc Natl Acad Sci U S A* 109, 2126-2131.
- Babkoff, H., Genser, S.G., Sing, H.C., Thorne, D.R., Hegge, F.W., 1985. The effects of progressive sleep loss on a lexical decision task: Response lapses and response accuracy. *Behavior Research Methods, Instruments, & Computers* 17, 614-622.
- Born, J., Feld, G.B., 2012. Sleep to upscale, sleep to downscale: balancing homeostasis and plasticity. *Neuron* 75, 933-935.
- Bosch, O.G., Rihm, J.S., Scheidegger, M., Landolt, H.P., Stampfli, P., Brakowski, J., Esposito, F., Rasch, B., Seifritz, E., 2013. Sleep deprivation increases dorsal nexus connectivity to the dorsolateral prefrontal cortex in humans. *Proc Natl Acad Sci U S A*.
- Braga, R.M., Sharp, D.J., Leeson, C., Wise, R.J., Leech, R., 2013. Echoes of the brain within default mode, association, and heteromodal cortices. *Journal Of Neuroscience* 33, 14031-14039.
- Chee, M.W., Choo, W.C., 2004. Functional imaging of working memory after 24 hr of total sleep deprivation. *Journal Of Neuroscience* 24, 4560-4567.
- Chee, M.W., Chuah, L.Y., Venkatraman, V., Chan, W.Y., Philip, P., Dinges, D.F., 2006. Functional imaging of working memory following normal sleep and after 24 and 35 h of sleep deprivation: Correlations of fronto-parietal activation with performance. *Neuroimage* 31, 419-428.
- Chee, M.W., Tan, J.C., 2010. Lapsing when sleep deprived: neural activation characteristics of resistant and vulnerable individuals. *Neuroimage* 51, 835-843.
- Chee, M.W., Tan, J.C., Parimal, S., Zagorodnov, V., 2010. Sleep deprivation and its effects on object-selective attention. *Neuroimage* 49, 1903-1910.
- Chee, M.W., Tan, J.C., Zheng, H., Parimal, S., Weissman, D.H., Zagorodnov, V., Dinges, D.F., 2008. Lapsing during sleep deprivation is associated with distributed changes in brain activation. *Journal Of Neuroscience* 28, 5519-5528.
- Chuah, L.Y., Dolcos, F., Chen, A.K., Zheng, H., Parimal, S., Chee, M.W., 2010. Sleep deprivation and interference by emotional distracters. *Sleep* 33, 1305-1313.
- De Havas, J.A., Parimal, S., Soon, C.S., Chee, M.W., 2012. Sleep deprivation reduces default mode network connectivity and anti-correlation during rest and task performance. *Neuroimage* 59, 1745-1751.
- Drummond, S.P., Brown, G.G., 2001. The effects of total sleep deprivation on cerebral responses to cognitive performance. *Neuropsychopharmacology* 25, S68-S73.
- Forster, K.I., Forster, J.C., 2003. DMDX: a windows display program with millisecond accuracy. *Behav Res Methods Instrum Comput* 35, 116-124.
- Fox, M.D., Snyder, A.Z., Vincent, J.L., Corbetta, M., Van Essen, D.C., Raichle, M.E., 2005. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A* 102, 9673-9678.
- Goel, N., Rao, H., Durmer, J.S., Dinges, D.F., 2009. Neurocognitive consequences of sleep deprivation. *Seminars In Neurology* 29, 320-339.
- Goldstein, A.N., Greer, S.M., Saletin, J.M., Harvey, A.G., Nitschke, J.B., Walker, M.P., 2013. Tired and apprehensive: anxiety amplifies the impact of sleep loss on aversive brain anticipation. *Journal Of Neuroscience* 33, 10607-10615.
- Greicius, M.D., Krasnow, B., Reiss, A.L., Menon, V., 2003. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc Natl Acad Sci U S A* 100, 253-258.
- Gujar, N., Yoo, S.S., Hu, P., Walker, M.P., 2010. The unrested resting brain: sleep deprivation alters activity within the default-mode network. *J Cogn Neurosci* 22, 1637-1648.
- Gusnard, D.A., Raichle, M.E., Raichle, M.E., 2001. Searching for a baseline: functional imaging and the resting human brain. *Nature Reviews Neuroscience* 2, 685-694.
- Heine, L., Soddu, A., Gomez, F., Vanhaudenhuyse, A., Tshibanda, L., Thonnard, M., Charland-Verville, V., Kirsch, M., Laureys, S., Demertzi, A., 2012. Resting state networks and consciousness: alterations of multiple resting state network connectivity in physiological, pharmacological, and pathological consciousness States. *Front Psychol* 3, 295.
- Kahnt, T., Chang, L.J., Park, S.Q., Heinzle, J., Haynes, J.D., 2012. Connectivity-based parcellation of the human orbitofrontal cortex. *Journal Of Neuroscience* 32, 6240-6250.

- Killgore, W.D., 2013. Self-reported sleep correlates with prefrontal-amygdala functional connectivity and emotional functioning. *Sleep* 36, 1597-1608.
- Killgore, W.D., Schwab, Z.J., Kipman, M., Deldunno, S.R., Weber, M., 2013. Insomnia-related complaints correlate with functional connectivity between sensory-motor regions. *Neuroreport* 24, 233-240.
- Kong, D., Soon, C.S., Chee, M.W., 2011. Reduced visual processing capacity in sleep deprived persons. *Neuroimage* 55, 629-634.
- Kong, D., Soon, C.S., Chee, M.W., 2012. Functional imaging correlates of impaired distractor suppression following sleep deprivation. *Neuroimage* 61, 50-55.
- L O Pez Zunini, R.I.O., Muller-Gass, A., Campbell, K., 2014. The effects of total sleep deprivation on semantic priming: Event-related potential evidence for automatic and controlled processing strategies. *Brain And Cognition* 84, 14-25.
- Larson-Prior, L.J., Power, J.D., Vincent, J.L., Nolan, T.S., Coalson, R.S., Zempel, J., Snyder, A.Z., Schlaggar, B.L., Raichle, M.E., Petersen, S.E., 2011. Modulation of the brain's functional network architecture in the transition from wake to sleep. *Progress In Brain Research* 193, 277-294.
- Li, Z., Kadivar, A., Pluta, J., Dunlop, J., Wang, Z., 2012. Test-retest stability analysis of resting brain activity revealed by blood oxygen level-dependent functional MRI. *Journal Of Magnetic Resonance Imaging* 36, 344-354.
- Libedinsky, C., Smith, D.V., Teng, C.S., Namburi, P., Chen, V.W., Huettel, S.A., Chee, M.W., 2011. Sleep deprivation alters valuation signals in the ventromedial prefrontal cortex. *Front Behav Neurosci* 5, 70.
- Lim, J., Choo, W.C., Chee, M.W., 2007. Reproducibility of changes in behaviour and fMRI activation associated with sleep deprivation in a working memory task. *Sleep* 30, 61-70.
- Lou, H.C., Luber, B., Crupain, M., Keenan, J.P., Nowak, M., Kjaer, T.W., Sackeim, H.A., Lisanby, S.H., 2004. Parietal cortex and representation of the mental Self. *Proc Natl Acad Sci U S A* 101, 6827-6832.
- Lythe, K.E., Williams, S.C., Anderson, C., Libri, V., Mehta, M.A., 2012. Frontal and parietal activity after sleep deprivation is dependent on task difficulty and can be predicted by the fMRI response after normal sleep. *Behavioural Brain Research* 233, 62-70.
- Mennes, M., Zuo, X.N., Kelly, C., Di Martino, A., Zang, Y.F., Biswal, B., Castellanos, F.X., Milham, M.P., 2011. Linking inter-individual differences in neural activation and behavior to intrinsic brain dynamics. *Neuroimage* 54, 2950-2959.
- Menz, M.M., Buchel, C., Peters, J., 2012. Sleep deprivation is associated with attenuated parametric valuation and control signals in the midbrain during value-based decision making. *Journal Of Neuroscience* 32, 6937-6946.
- Minkel, J.D., McNealy, K., Gianaros, P.J., Drabant, E.M., Gross, J.J., Manuck, S.B., Hariri, A.R., 2012. Sleep quality and neural circuit function supporting emotion regulation. *Biol Mood Anxiety Disord* 2, 22.
- Mu, Q., Mishory, A., Johnson, K.A., Nahas, Z., Kozel, F.A., Yamanaka, K., Bohning, D.E., George, M.S., 2005. Decreased brain activation during a working memory task at rested baseline is associated with vulnerability to sleep deprivation. *Sleep* 28, 433-446.
- Mu, Q., Nahas, Z., Johnson, K.A., Yamanaka, K., Mishory, A., Koola, J., Hill, S., Horner, M.D., Bohning, D.E., George, M.S., 2005. Decreased cortical response to verbal working memory following sleep deprivation. *Sleep* 28, 55-67.
- Mullin, B.C., Phillips, M.L., Siegle, G.J., Buysse, D.J., Forbes, E.E., Franzen, P.L., 2013. Sleep deprivation amplifies striatal activation to monetary reward. *Psychological Medicine* 43, 2215-2225.
- Muto, V., Shaffii-le, B.A., Matarazzo, L., Foret, A., Mascetti, L., Jaspar, M., Vandewalle, G., Phillips, C., Degueldre, C., Baletau, E., Luxen, A., Collette, F., Maquet, P., 2012. Influence of acute sleep loss on the neural correlates of alerting, orientating and executive attention components. *Journal Of Sleep Research* 21, 648-658.
- Niendam, T.A., Laird, A.R., Ray, K.L., Dean, Y.M., Glahn, D.C., Carter, C.S., 2012. Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. *Cogn Affect Behav Neurosci* 12, 241-268.
- Oldfield, R.C., 1971. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9, 97-113.
- Parsons, C.E., Stark, E.A., Young, K.S., Stein, A., Kringelbach, M.L., 2013. Understanding the human parental brain: A critical role of the orbitofrontal cortex. *Soc Neurosci* 8, 525-543.
- Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 59, 2142-2154.
- Raichle, M.E., 2010. Two views of brain function. *Trends In Cognitive Sciences* 14, 180-190.
- Raichle, M.E., 2011. The restless brain. *Brain Connect* 1, 3-12.

Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L., 2001. A default mode of brain function. *Proc Natl Acad Sci U S A* 98, 676-682.
 Samann, P.G., Tully, C., Spoormaker, V.I., Wetter, T.C., Holsboer, F., Wehrle, R., Czisch, M., 2010. Increased sleep pressure reduces resting state functional connectivity. *MAGMA* 23, 375-389.
 Shao, Y., Wang, L., Ye, E., Jin, X., Ni, W., Yang, Y., Wen, B., Hu, D., Yang, Z., 2013. Decreased thalamocortical functional connectivity after 36 hours of total sleep deprivation: evidence from resting state FMRI. *PLoS One* 8, e78830.
 Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins, K.E., Toro, R., Laird, A.R., Beckmann, C.F., 2009. Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci U S A* 106, 13040-13045.
 Sterpenich, V., Albouy, G., Darsaud, A., Schmidt, C., Vandewalle, G., Dang, V.T., Desseilles, M., Phillips, C., Degueldre, C., Balteau, E., Collette, F., Luxen, A., Maquet, P., 2009. Sleep promotes the neural reorganization of remote emotional memory. *Journal Of Neuroscience* 29, 5143-5152.
 Thomas, M., Sing, H., Belenky, G., Holcomb, H., Mayberg, H., Dannals, R., Wagner, H., Thorne, D., Popp, K., Rowland, L., Welsh, A., Balwinski, S., Redmond, D., 2000. Neural basis of alertness and cognitive performance impairments during sleepiness. I. Effects of 24 h of sleep deprivation on waking human regional brain activity. *Journal Of Sleep Research* 9, 335-352.
 Tononi, G., Cirelli, C., 2003. Sleep and synaptic homeostasis: a hypothesis. *Brain Research Bulletin* 62, 143-150.
 Tononi, G., Cirelli, C., 2006. Sleep function and synaptic homeostasis. *Sleep Medicine Reviews* 10, 49-62.
 Turner, J.A., Chen, H., Mathalon, D.H., Allen, E.A., Mayer, A.R., Abbott, C.C., Calhoun, V.D., Bustillo, J., 2012. Reliability of the amplitude of low-frequency fluctuations in resting state fMRI in chronic schizophrenia. *Psychiatry Res* 201, 253-255.
 Uddin, L.Q., Kelly, A.M., Biswal, B.B., Castellanos, F.X., Milham, M.P., 2009. Functional connectivity of default mode network components: correlation, anticorrelation, and causality. *Human Brain Mapping* 30, 625-637.
 Van Dijk, K.R., Sabuncu, M.R., Buckner, R.L., 2012. The influence of head motion on intrinsic functional connectivity MRI. *Neuroimage* 59, 431-438.
 Van Dongen, H.P., 2005. Brain activation patterns and individual differences in working memory impairment during sleep deprivation. *Sleep* 28, 386-388.
 Yan, C.G., Cheung, B., Kelly, C., Colcombe, S., Craddock, R.C., Di Martino, A., Li, Q., Zuo, X.N., Castellanos, F.X., Milham, M.P., 2013. A comprehensive assessment of regional variation in the impact of head micromovements on functional connectomics. *Neuroimage* 76, 183-201.
 Yan, C.G., Craddock, R.C., Zuo, X.N., Zang, Y.F., Milham, M.P., 2013. Standardizing the intrinsic brain: towards robust measurement of inter-individual variation in 1000 functional connectomes. *Neuroimage* 80, 246-262.
 Zald, D.H., McHugo, M., Ray, K.L., Glahn, D.C., Eickhoff, S.B., Laird, A.R., 2012. Meta-Analytic Connectivity Modeling Reveals Differential Functional Connectivity of the Medial and Lateral Orbitofrontal Cortex. *Cerebral Cortex*.
 Zang, Y.F., He, Y., Zhu, C.Z., Cao, Q.J., Sui, M.Q., Liang, M., Tian, L.X., Jiang, T.Z., Wang, Y.F., 2007. Altered baseline brain activity in children with ADHD revealed by resting-state functional MRI. *Brain Dev* 29, 83-91.
 Zou, Q., Ross, T.J., Gu, H., Geng, X., Zuo, X.N., Hong, L.E., Gao, J.H., Stein, E.A., Zang, Y.F., Yang, Y., 2013. Intrinsic resting-state activity predicts working memory brain activation and behavioral performance. *Human Brain Mapping* 34, 3204-3215.
 Zuo, X.N., Di Martino, A., Kelly, C., Shehzad, Z.E., Gee, D.G., Klein, D.F., Castellanos, F.X., Biswal, B.B., Milham, M.P., 2010. The oscillating brain: complex and reliable. *Neuroimage* 49, 1432-1445.

Table 1. Brain areas of ALFF differences between the RW and SD group. Coordinates x, y, z (mm) are given in standard stereotactic MNI space. All regions listed are statistically significant at the $p < 0.05$, AlphaSim corrected. L: left; R: right.

SD1:

Brain regions	MNI coordinates			BA	L/R	Voxels	T values
	x	y	z				
Superior Frontal Gyrus	-12	66	6	10	L	202	-4.27
Middle Frontal Gyrus	45	27	30	46	R	47	-3.62
Middle Frontal Gyrus	-45	39	-6	46/47	L	47	-4.03
Inferior Frontal Gyrus	54	18	0	22/47	R	82	-4.01
Middle Occipital Gyrus	-51	-66	-15	19/37	L	653	4.66
Middle Occipital Gyrus	42	-60	-9	19	R	109	4.35
Middle Occipital Gyrus	-12	-72	0	18/19	L	101	3.99
Thalamus	21	-15	12	-	R	98	4.37
Thalamus	-18	-18	3	-	L	40	4.19
Inferior Parietal Lobule	45	-60	33	39/40	R	148	-5.48
Precuneus/ Posterior Cingulate	3	-63	30	31/7	R	70	-5.58
Precentral Gyrus	-36	-9	60	1/2/3/4/6	L	738	7.32
Precentral Gyrus	-54	-3	36	3/4/6	L	41	4.14

SD2:

Brain regions	MNI coordinates			BA	L/R	Voxels	T values
	x	y	z				
Medial Frontal Gyrus	15	66	-3	11/10	R	54	-5.22
Superior Frontal Gyrus	-42	48	0	10/46	L	39	-4.62
Middle Frontal Gyrus	-51	33	27	9/46	L	40	-5.01
Inferior Frontal Gyrus	-51	36	0	10/47	L	62	-4.62
Insula	-33	-18	21	13	L	378	8.99
Insula	36	-15	21	13	R	56	4.67
Post Cingulate Gyrus	0	-33	33	23/31	L	30	-4.51
Inferior Parietal Lobule	66	-48	24	39/40	R	156	-6.09
Precentral Gyrus	36	-24	36	2/3/4/6	R	319	6.51
Precentral Gyrus	-36	-12	57	1/2/3/4/6	L	652	10.06
Middle Occipital Gyrus	-39	-87	0	18/19	L	413	6.79
Medial Frontal Gyrus	15	66	-3	11/10	R	54	-5.22
Superior Frontal Gyrus	-42	48	0	11/46	L	39	-4.62

Figure Legends




Figure 1. Effects of 24h sleep deprivation on RT for semantic tasks. Y-axes show the mean (± 2 sd) time of RT for each group. X-axes indicate the group. Significantly different (* $p < 0.05$, ** $p < 0.01$) groups are as indicated.




Figure 2: Group ALFF differences in the two SD sessions. The effects are significant at $p < 0.05$, AlphaSim corrected. Cool color indicates that the SD group had decreased ALFF compared with the controls and the hot color indicates the opposite. The results were visualized with the BrainNet Viewer (<http://www.nitrc.org/projects/bnv/>).




Figure 3: Overlap of SD1 and SD2 compared to control differences. 2 SD sessions shared the same areas are shown in brown and blue. **Voxel-wise ICC.** The intra-subject test-retest reliability for two SD sessions, the regions with high reliability ($ICC > 0.5$) are shown. The results were visualized with the BrainNet Viewer (<http://www.nitrc.org/projects/bnv/>).




Figure 4: The correlation between RTs and ALFF change in the right IPL in the sleep deprivation group. The greater the prolongations in RT from rest to sleep deprivation, the greater the decrease in ALFF in the right IPL. The results were visualized with the BrainNet Viewer (<http://www.nitrc.org/projects/bnv/>).